

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP Indian Journal of Anatomy and Surgery of Head, Neck and Brain

Journal homepage: <https://www.ijashnb.org/>

## Case Report

# Aphasia, apraxia, and dysphagia following viper envenomation: A single case report

R Johnsi Rani<sup>1</sup>, V Jaya<sup>1</sup>, V Monish<sup>2,\*</sup>

<sup>1</sup>Institute of Speech & Hearing, Madras Medical College, Chennai, Tamil Nadu, India

<sup>2</sup>Dept. of Speech, Hearing & Communication, National Institute for Empowerment of Persons with Multiple Disabilities (Divyangjan), Chennai, Tamil Nadu, India



### ARTICLE INFO

#### Article history:

Received 16-03-2022

Accepted 19-03-2022

Available online 02-04-2022

#### Keywords:

Viper Envenomation

Aphasia

Apraxia

Dysphagia

### ABSTRACT

**Objective:** Aim of the work was to highlight unusual complications such as aphasia, apraxia and dysphagia following viper bite.

**Method:** In this case report, a 55 years old male who had developed Cerebrovascular Accident (CVA) and Right hemiparesis after an hour of saw-scaled viper bite on his right foot was described. His MRI report revealed chronic infarct in bilateral fronto parietal lobe. Difficulty in speaking and swallowing were observed and the patient was under nasogastric feeding. For assessing speech and language skills, Western Aphasia Battery and Apraxia Battery for Adults-2 was used. Manipal Manual for Swallowing Assessment was used for assessing the swallowing skills.

**Results:** The assessment results indicated that the patient had Broca's aphasia with Non-Verbal oral apraxia and oropharyngeal dysphagia. Improvement was observed in swallowing and verbal communication skills following speech-language intervention.

**Conclusion:** Aphasia, apraxia and dysphagia were unusual complications following viper bite. Therefore, it is important for a speech language pathologist to have knowledge regarding various causes of stroke for better speech, language and swallowing management

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: [reprint@ipinnovative.com](mailto:reprint@ipinnovative.com)

## 1. Introduction

Snake envenomation is quite usual problem in tropical countries which affect nearly six million people each year, with approximately 10% of these fatalities happening in India.<sup>1</sup> Neurological deficit following viper envenomation is not uncommon. Generally it manifests as intracerebral hemorrhage or subarachnoid hemorrhage.<sup>2</sup> In India, the leading cause of fatal snake bite are viperidae species which consists of Russell's viper (*Daboia russelli*) and saw scaled viper (*Echis carinatus*).<sup>3-5</sup> Snake envenomation is a complicated phenomenon. The manifestation of snake envenomation includes hematological, cardiac,

renal and neurological abnormality. Though ischemic manifestations such as, acute myocardial infarction or cerebral infarction is quite unusual.<sup>6,7</sup> Snake envenomation leads to a diverse range of clinical effects, ranging from minimal envenomation like itching, pain, puncture wounds and swelling to highly fatal manifestations such as bleeding, shock, hypotension, and renal failure, specifically with viper bites, and respiratory failure and sudden flaccid paralysis in neurotoxic bites.<sup>8</sup> Saw scaled viper envenomation (*Echis carinatus*) leads to local swelling and severe necrosis of tissue. This manifestation includes hemorrhage in various parts of the body such as gums, nasopharynx, gastrointestinal tract, central nervous system and urinary tract. The neurological traits includes

\* Corresponding author.

E-mail address: [monishvasudevan@gmail.com](mailto:monishvasudevan@gmail.com) (V Monish).

drowsiness, convulsions and confusions.<sup>9</sup> There is a dearth of research studies reporting specifically on the speech, language and swallowing deficits that occur following snake bite, hence in this case report we present unusual complications such as aphasia, apraxia and dysphagia following viper bite.

## 2. Case Report

A case of 55 years old male had developed Cerebrovascular Accident (CVA) and Right hemiparesis after an hour of saw-scaled viper bite on his right foot. His MRI report revealed chronic infarct in bilateral fronto parietal lobe. He had difficulty in speaking and swallowing and the patient was under nasogastric feeding. He had no other history of risk factors such as diabetes mellitus, hypertension, smoking, alcohol consumption. For speech and swallowing evaluation, the patient was referred to Institute of Speech and Hearing, Madras Medical College, Chennai, Tamil Nadu.

### 2.1. Speech, language and swallowing evaluation

A comprehensive assessment was carried out by the speech language pathologist. The native language of the patient was Tamil. He was right handed, monolingual and completed 10<sup>th</sup> standard. His hearing skills were normal. During the evaluation, he was conscious, alert and oriented though emotionally unstable. On speech and language evaluation, he had non-fluent speech, impairment in naming, reading, writing, repetition with good auditory and visual comprehension. His mode of communication was Non-Verbal, through gestures and pointing. Oral Peripheral Mechanism Examination was carried out. Western Aphasia Battery<sup>10</sup> is an accepted method for evaluating language skills in many neurological conditions such as stroke.<sup>11,12</sup> In Western Aphasia Battery specifically six subscales of the test which includes spontaneous speech, comprehension, naming, repetition, reading and writing was administered. As a measure of apraxia, subscores from the Apraxia Battery for Adults-2<sup>13</sup> (ABA-2) was used. For swallowing evaluation, Manipal Manual for Swallowing Assessment<sup>14</sup> (MMSA) was used.

## 3. Results

Oral Peripheral Mechanism Examination revealed lips deviated towards left side with reduced lip closure, poor lip seal; tongue deviated towards right side and restricted lingual movements.

On WAB,<sup>10</sup> the subtest scores were 0 for spontaneous speech, 8.2 for comprehension, 0 for naming, repetition, reading and writing revealing Broca's aphasia.

On Apraxia Battery for Adults (ABA) indicated Non-Verbal oral apraxia.

MMSA scores on motor and sensory assessment were 104 and on phases of swallowing were 19 revealing Dysphagia in oral and pharyngeal phase. He had delayed laryngeal elevation while swallowing thick and thin liquids. Aspiration and cough with expectoration was observed while swallowing saliva and thin liquids. Diminished gag reflex was observed. He was diagnosed as Broca's aphasia with Non-Verbal oral apraxia and oropharyngeal dysphagia.

## 4. Intervention

The patient has been undergoing speech, language and swallowing intervention for the past one month. Oromotor exercises such as lip stretching, lip closure exercises were given to improve lip seal and labial range of motion. Tongue stretching and strengthening exercises were given to elicit tongue movements and range of motion. Vegetative skills such as blowing and sucking were practiced.

Swallowing facilitation techniques were carried out to manifest adequate swallowing. Diet modification was used starting from spoon thick consistency to thin liquids. Head extension technique was used to drain the food from the oral cavity to the pharynx and oesophagus through gravity. Effortful swallow was done to increase the posterior movement of tongue base during the pharyngeal swallow to improve the clearance of bolus from vallecula. Stimulation was given using food of different tastes and temperatures alternatively to enhance the taste sensation and to elicit the gag reflex.

After a month of follow up, he was able to swallow thick, thin liquids and semi-solids with no aspiration. At present, he is able to maintain good lip seal and tongue movements such as protrusion, retraction and elevation were elicited. Blowing and sucking skills were also established. His MMSA scores were reduced from 104 on sensory and motor assessment to 93 and on phases of swallowing from 19 to 10. Even though intervention was started one year after the snake bite, he has shown marked improvement in oromotor and swallowing skills.

With speech and language intervention, improvements were observed in verbal communication skills. After a month of follow up, he was also able to comprehend complex sentences. He can speak in sentence but grammatical errors were observed.

## 5. Discussion

Serious neurological abnormalities, including stroke and muscle paralysis, are associated with the toxic effects of the venom. The venom consists of a complex mixture of toxins influencing the neuromuscular transmission, the coagulation cascade, or both.<sup>15</sup> The rich components present in the complex toxin of viper snake venom dominantly affect the hemostatic mechanism.<sup>16</sup> Characteristics of viper bite depend on the envenomation severity and

bleeding manifestations are frequently encountered in clinical practice.<sup>17</sup> In large doses, viper snake venom can lead to massive intravascular coagulation which cause occlusion of small and even large vessel resulting in cerebral infraction.<sup>18</sup> Bashir and Jinkins (1985)<sup>19</sup> reported a patient in whom envenomation with *Echis carinatus* (saw scaled viper) resulted in cerebral infraction. That patient developed hemiplegia and aphasia (non-fluent aphasia). Polo, Alvarez de Arcaya, Cid, Berciano (2002)<sup>20</sup> reported a farmer who developed sensory aphasia following viper bite. They also mentioned that the patient was eventually improved with treatment conservatively and residual expressive aphasia was the patient's only sequel when last seen. Numeric, Moravie, Didier. Chatot-Henry D, Cirille (2002)<sup>21</sup> reported a patient who developed multiple cerebral infarctions after snake envenomation. He had Partial wernicke's aphasia (including difficulty in repetition of spoken words and also difficulty in simple straightforward commands). Narang, Paleti, Asad, and Samina (2009)<sup>5</sup> reported right hemiplegia and expressive aphasia in a patient following a Russell's viper bite. They also mentioned improvement in motor power and speech in that patient. Chandrashekar Anikethana, and Kalinga (2012)<sup>22</sup> described a patient who developed right hemiplegia with motor aphasia caused due to infraction following viper bite. Also, they mentioned that the patient had improved with nominal aphasia. There were other studies which reported expressive aphasia,<sup>23</sup> motor aphasia,<sup>24</sup> mixed aphasia<sup>25</sup> following snake bite.

In our patient he was previously healthy with no significant premorbid history, developed broca's aphasia as a result of cerebral infraction following snake envenomation. To our knowledge, this is the first article to report non-verbal oral apraxia associated with broca's aphasia following snake envenomation.

Dysphagia is one of the neurotoxic features following snake bite.<sup>26</sup> Bernheim, Lorenzetti, Licht, Markwalder and Schneemann (2001)<sup>27</sup> reported three cases of severe neurotoxicity following snake bite. One of the three patients exhibited dysphagia. Seneviratne and Dissanayake (2002)<sup>28</sup> mentioned that dysphagia is one of the neurological manifestation after snake bite. Similarly, our patient developed oropharyngeal dysphagia following snake envenomation. With swallowing therapy, the patient's swallowing ability was improved, and he was able to continue oral feeding.

## 6. Conclusion

In snake envenomation, the mechanism through which the cerebral infraction happens can be multi-factorial. The Toxic snake venom contains anti-coagulant and pro-coagulant properties.<sup>29</sup> It can be a possible cause for stroke resulting in aphasia with apraxia as well as dysphagia. Therefore it is important for a speech language pathologist to have knowledge regarding various causes of stroke for

better speech, language and swallowing management. The speech, language and swallowing skills have been improved after proper intervention. If the intervention had started earlier, improvement would have been much better.

## 7. Source of Funding

None.

## 8. Conflict of Interest

The author declares that there is no conflict of interest.

## References

1. Kasturiratne A, Wickremasinghe AR, De Silva N, Gunawardena NK, Pathmeswaran A, Premaratna R, et al. The global burden of snakebite: A literature analysis and modeling based on regional estimates of envenoming and deaths. *PLoS Med.* 2008;5(11):218. doi:10.1371/journal.pmed.0050218.
2. Murthy JM, Kishore LT, Naidu KS. Cerebral infarction after envenomation by viper. *J Comput Assist Tomogra.* 1997;21(1):35–7. doi:10.1097/00004728-199701000-00007.
3. David AW. Guidelines for the clinical management of snake-bites in South-East Asia region.; 2005. Available from: <https://apps.who.int/iris/handle/10665/205171>.
4. Brunda G, Sashidhar RB. Epidemiological profile of snake-bite cases Andhra Pradesh using immunoanalytical approach. *Indian J Med Res.* 2007;125(5):661–8. [17642502].
5. Narang SK, Paleti S, Asad MA, Samina T. Acute ischemic infarct in the middle cerebral artery territory following a Russell's viper bite. *Neurol India.* 2009;57(4):479–80. doi:10.4103/0028-3886.55594.
6. Hsaini Y, Satte A, Balkhi H, Karouache A, Bourezza A. Stroke following a viper bite. *Ann Fr Anesth Reanim.* 2010;29(4):315–6.
7. Aravanis C, Ioannidis PJ, Ktenas J. Acute myocardial infarction and cerebrovascular accident in a young girl after a viper bite. *Br Heart J.* 1982;47(5):500–3. doi:10.1136/hrt.47.5.500.
8. Jones AL, Haslett C, Chilvers ER, Boon N, Colledge NR, Hunter JAA, et al. Davidson's Principles and Practice of Medicine; 2002. p. 166–84.
9. Warrell DA, Nmed D, Greenwood BM, Ormerod LD, Pope HM, Watkins BJ, et al. Poisoning by bites of the saw-scaled or carpet viper (*Echis carinatus*) in Nigeria. *Q J Med.* 1977;46(181):33–62. [866568].
10. Kertesz A. Western Aphasia Battery. New York: Grune & Stratton; 1982. doi:https://doi.org/10.1007/978-0-387-79948-3\_937.
11. Mamikonyan E, Moberg PJ, Siderowf A, Duda JE, Have TT, Hurtig HI, et al. Mild cognitive impairment is common in Parkinson's disease patients with normal Mini-Mental State Examination (MMSE) scores. *Parkinsonism Relat Disord.* 2009;15(3):226–31. doi:10.1016/j.parkreldis.2008.05.006.
12. Janvin CC, Larsen JP, Aarsland D, Hugdahl K. Subtypes of mild cognitive impairment in Parkinson's disease: progression to dementia. *Mov Disord.* 2006;21(9):1343–9. doi:10.1002/mds.20974.
13. Dabul B. Apraxia Battery for Adults-Second Edition. 2000;.
14. Balasubramaniam RK, Bhat JS. Manipal manual for swallowing assessment. Manipal: Manipal University Press; 2012.
15. Brutto OHD, Brutto VJD. Neurological complications of venomous snake bites: a review. *Acta Neurologica Scandinavica.* 2012;125(6):363–72. doi:10.1111/j.1600-0404.2011.01593.x.
16. Marsh NA. Snake venom affecting haemostatic mechanism - A consideration of their mechanism, practical applications and biological significance. *Blood Coagul Fibrinolysis.* 1994;5(3):399–410. [8075311].
17. Chugh KS. Snake bite induced acute renal failure in India. *Kidney Int.* 1989;35(3):891–907. doi:10.1038/ki.1989.70..
18. Schwartzman RJ, Hill JB. Neurologic complications of disseminated intravascular coagulation. *Neurology.* 1982;32(8):791–7.

- doi:10.1212/wnl.32.8.791.
19. Bashir R, Jinkins J. Cerebral infarction in a young female following snake bite. *Stroke*. 1985;16(2):328–30. doi:10.1161/01.str.16.2.328.
  20. Polo JM, De Arcaya AA, Cid C, Berciano J. Aphasia in a farmer following viper bite. *Lancet*. 2002;359(9324):2164. doi:10.1016/s0140-6736(02)09094-3.
  21. Numeric P, Moravie V, Didier M, Chatot-Henry D, Cirille S. Multiple cerebral infarctions following a snakebite by *Bothrops caribbaeus*. *Am J Trop Med Hyg*. 2002;67(3):287–8. doi:10.4269/ajtmh.2002.67.287.
  22. Anikethana GVC, Kalinga BE. *Viper Bite Presenting as Acute Ischemic Stroke*. 2012;3(11).
  23. Pothukuchi VK, Chepuri VR, Natta K, Madigani N, Kumar A. A rare case report of Russell's viper snakebite with ischemic stroke. *Hong Kong Journal of Emergency Medicine*. 2017;25(2):95–7. doi:https://doi.org/10.1177/1024907917735071.
  24. Ajit D, Kumar SG. *Acute Cerebral Infarct on Evolution in Middle Cerebral Artery Following Viper Snake Bite*. 2016;5(1). doi:10.21275/v5i1.nov153081.
  25. Zeng X, Hu J, Liang X, Wu Y, Yan M, Zhu M, et al. Acute cerebral infarction following a *Trimeresurus stejnegeri* snakebite: A case report. *Medicine*. 2019;98(23):e15684. doi:10.1097/MD.00000000000015684.
  26. Amin AR, Mamun SMH, Chowdhury NH, Rahman M, Ali M. Neurotoxic manifestation of snake bite in Bangladesh. *J Clin Toxicol*. 2014;4(185):2161–0495. doi:10.4172/2161-0495.1000185.
  27. Bernheim A, Lorenzetti E, Licht A, Markwalder K, Schneemann M. Three cases of severe neurotoxicity after cobra-bite (*Naja kaouthia*). *Swiss Med Wkly*. 2001;131(15-16):227–8. [11400547].
  28. Seneviratne U, Dissanayake S. Neurological manifestations of snake bite in Sri Lanka. *Journal of postgraduate medicine*. 2002;48(4):275–8. discussion 278-9. [12571382].
  29. Paul G, Paul BS, Puri S. Snake bite and stroke: Our experience of two cases. Indian journal of critical care medicine: peer-reviewed, official publication of Indian Society of. *Critical Care Medicine*. 2014;18(4):257–8. doi:10.4103/0972-5229.130585.

## Author biography

**R Johnsi Rani**, Faculty

**V Jaya**, Faculty

**V Monish**, Faculty

**Cite this article:** Rani RJ, V Jaya, V Monish. Aphasia, apraxia, and dysphagia following viper envenomation: A single case report. *IP Indian J Anat Surg Head, Neck Brain* 2022;8(1):26-29.