

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP Indian Journal of Anatomy and Surgery of Head, Neck and Brain

Journal homepage: <https://www.ijashnb.org/>

Original Research Article

Analysis of anatomy of head and neck: A study through empiricals part- II

V Rama Raju^{1,*}

¹CMR College of Engineering & Technology, Affili.: Jawaharlal Nehru Technological University JNTU, Hyderabad, Telangana, India



ARTICLE INFO

Article history:

Received 15-01-2023

Accepted 25-01-2023

Available online 20-02-2023

Keywords:

Sterno-cleido-mastoid

Surgical-operations

ABSTRACT

In this study we introduce new hypothesis into the anatomical-structure, the head, and neck, neuro-embryology, the nerves and muscles, followed by the physiologic-variants, operational / surgical implications and considerations and then lastly clinical significance of the study. The wound/damage to the cervical altruistic ganglia can elicit the syndrome-of-Horner, which is ipsilateral ptosis, miosis, and also facial scrub—anhidrosis. The torticollis-transpires as sterno-cleido-mastoid (SCM) muscle-tone abbreviates spasmodically, generating the fluctuating of the neck opposite to aomolous muscle-tome cross or control leads to in biotic if the sterno-cleido-mastoid is incapacitated, also that might be leading to fibrosis as well as margarine of ‘muscle-fiberes’. And as well turn out as a consequence of enhanced muscle - tone, and/or dystonia muscle spasms and spasms, lessening from the poignant and sensitive nervousness stress and strain and tension also, and/or unanticipated behaviors. Though, even though, whilst evaluating the traumatic injuries or wounds, it is so crucial to omit or inhibit certain carotid monitorings as well as distinguishing in the interior of the neck.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

The human-head stands on top of the body-vertebra together with the brain-cranium (the skull) linking and join up at initial cervical-vertebra on the “atlas” referred to as. The thin portion of the head—neck shapes the best upper portion of the axial—skeleton as well as it is also build with the brains=cranium (the skull), hyoid-bone followed by the spine-cervical.¹⁻³ It is the structural(anatomical) component which consists of the brain-cranium(skull), hyoid—bone and also the vertebrae(cervical). Here, skull is referred to as mutually indicating mandible (which is a lower—jaw of the bone) plus brain-cranium (which is an higher-part of the cranium-head which stores the human-brain).

The human-neck (shaft, or neckline) anatomical-structures are spread in the interior four-spot partitions.⁴⁻⁷

* Corresponding author.

E-mail address: drvenkateshwararr@gmail.com (V. R. Raju).

1. “Vertebral—section” which includes cervical-vertebra, through the cartilaginous-disc amid each, and everybody-vertebra. However, it is defined the orientation of the body—vertebrae derives the forms (shaping) of the neck of the human.⁵
2. As and when the vertebra tied the “spinal—canal” (canal of the spine), the “cervical—section” which belongs to “spinal—cord” was well detected surrounded by the neck opening.
3. “Visceral—section” that holds the throat(trachea), larynx(related to voice), throat(pharynx), thyroid followed by para thyroid secretors(glands, secretory/ or secretory organ).
4. “Vascular—section” is come together as well as comprises two carotids—sheath uncovered lying on top of each and every-part of the throat(‘trachea’). All the carotids—sheath consists of vagus—nerve, normal

carotid—vein(‘artery’ and/or route) as well as the interior (inner) jugular-artery(‘vein’).

Above and beyond and also likewise, the registered anatomical—structures, the neckline comprises lymph’s, i.e.,(cervical—lymph’s) joints (nodes) or connections which encompass the plasma (life-blood)-vessels.^{6–10}

2. Plasma—Supply, and lymphatics

Most significant plasma—vessels of human—neck are restricted and also restricted contained by the surrounded by the sheath of the carotids which are normal carotid—arteries, inner carotids—arteries, followed by the interior vena jugularis jugular—veins.

The “right-side” normal carotidarteries derives as of brachio-cephalic-arteries, whereas left-side normalcarotid arteries derives right away from ‘aortic-archies’. The usual carotids separates hooked on the interior, the inner-carotids followed by the outer carotids-arteries maximum to the extent of the exceptional section of the “thyroid-cartilage.^{1,7}” The exterior carotidarteries egresses the sheaths of carotids plus provides the cursory and/or trivial characteristic of the side (face-look) as well as components of neckline and that yields exceptions—thyroid arteries, linguisticartery (lingual), facial-artery, plus occipital—arteries. The inner carotidartery remains in to lay-bone all through across the canals of the carotids plus connects the circle-of-Willis to provide the ocular(ophthalmic-opthamology) blood vessel, frontal intellectual (cerebral) arteries, also central intellectual arteries.^{1,7}

Numerous lymph connections are within the neck-of-the head, through bulk positioned or situated adjacent to the path of the interior vena jugularis-veins. The horizontal head-neck lymph-connections occur within frontal plus backside shackles over every-side section of head-neck, adjacent to as well as strongly connected through inner jugulars—veins.

2.1. Neuro embryology

The embryology which is a field (branch) of science which is associated with the development, evolution, as well as advancement of nucleus which is dealing through the prenatal—phase of growth opening as of creation of gametes, insemination, creation-of-zygote(fertilized-ovum), growth-of-embryo as well as the fetus to the natal of a novel/new-fangled novel separate. There are mainly two fundamental procedures concerned are development as well as diversity that leads to the creation of different tissues also tissues and structural-tissues within the body-vertebra dedicated to stage to accomplish particular tasks and purposes. This neuro embryology field is is linked to the growth of the CNS which consists of the human brain as well as the spinal plus the PNS – the prephiral nervous-

system which consists of the spinal/vertebral, cranial, as well as “autonomic-nerves” within the human-body’. Such that these soft—tissues build as of neural/ or neuronal-tube also neural/neuronal-growth-cells.

2.2. Nerves

The human head-neckline is house of the variety of anatomical structures of the nervous—system.

The cervical—gangliaon is the threesome triangle of sensitive nervous-system(9NS) gangliaon which sit together with the vertebral/spinal row/column. The exceptional cervical-ganglia lie down at the C 2 / C 3 entomb or intomb vertebral/spinal height, whilst the central cervical—ganglia remains rests at the C 6 / C 7 lay to rest spinal/vertebral degree point. The mediocre(inferior) cervic ganglia is combined through a few thoracic—ganglia to build the “stellate—ganglia” at the C 7 / T 1 entomb spinal/vertebral intensity-point.^{8–10}

The cranial-nerves, i.e.,X.I (spinal/vertebra-attachment-nerve) has a cranial-source as well as the spinal/vertebral origin. The cranial foundation leaves the vena jugularis for a men also connects the ‘vagusnerves. That traverses the subsequent cervical-trio threesome-triangle at the bottom or subsequent “digastric-muscle” as well as remains substandard while waiting for it innervate the sterno-cleido-mastoid (SCM)muscle. Later that maintains externally to the “levator-scapulae” LSM(muscle and also completes in the musculus-trapezius.⁸




Fig. 1: Veins (arteries) of human—head plus neck. https://en.wikipedia.org/wiki/Inferior_tympanic_artery

2.3. Muscles—nerves

The platysma is one of the slender-muscle which stretches as of the superior chest-pectus(thorax) towards the cheek-line (brass) also slighter-perimeter. Such that it works to attraction the dominant rim substandard, as well as the strained the neckline externally, also gets nerve source as of the ‘cervical-branch’ of the “facial-nerves”.¹¹

The sterno- cleido- mastoid (SCM) muscle has had two heads of muscles which derive as of the sternal manubriums as well as ‘median-clavicle’. These heads of muscles combine plus include at the mastoid-process process of the sequential/temporal-bone as well as exceptional nuchalneckline. Such that which works to turn around the top to the opposing position which is tightening. So, its innervation is through the C N - X I.^{12–14}

The suprahyoid trapezius muscles comprise of the digastric, mylohyoid, as well as geniohyoid—muscles’. Such that these muscle-tones ascribe attribute to the bone of-hyoid as well as segments of the jaw bones, with the exception of for the “digastric—muscle.”

The digastrics muscles that has two tummies, one is the latter of which connects to the mastoidal-processes of the terrestrial-bone’. Innervated of the genio-hyoid’ is through the C N X-II, which is also referred to as the “hypo-glossal-nerve”. The frontal tummy of the digastrics- muscles as well as the mylohyoid pick up nerve source as of the mylohyoid-nerve’, a offshoot of the inframaxillary branch of the C N V. The facial scrub nerve enervated the digastrics-muscle rear end abdomen. Such that their work is to improve the “hyoids-bones”.¹

2.4. Physiological-variations

The then peripheral carotid-arteries typically yields off 3 frontal sections. On or after lower to excellent, they are the greater thyroids arteries, the linguistic, i.e., lingual-arteries artery’s, as well as the facial’ scrub artery’s.

The exceptional thyroid-artery’s pass through substandard to the ‘thyroid-gland’ which also yieldsoff the sternomastoid(SC) artery’s. Sporadically, and intermittently the SCartery will occur immediately as of the peripheral “carotid-artery’s”.^{14–18}

The facial scrub artery’s typically leaves the peripheral carotid-artery’s higher than the beyond the higher-horn’ of the hyoid also which pass through commandingly at the back of the sub mandi bular-glands.

3. Operational Matters and Consequences

In the course of numerous shoulder joint operations (surgical-operations), the “nerve-blocks” are employed to reduce discomfort for the duration of throughout the course of the initial twenty-four hours, i.e., one complete day of operation, that enhances the subject who is a patient patient fulfillment, as well as it reduces the demand for

“opioid-discomfort” alleviation. Such that these “nerve-blocks”, nerve-jumps are frequently performed within the ultra sound management as well as are transferred on to the “brachial-plexus’ muscles” right away directly as they leave the frontal as well as the central “scalene-muscles” within the neckline. However, there are several indeed many and also more cervical-cell “blocks/units” employed in to the surgical operation.^{15–22}

4. Conclusions

Approximately circa ~40% of head followed by the neck surgeries are linked through the post op (post operative) disease, that frequently and more often happens within a month following the operational-surgery. Therefore, further study is followed. In the coming research study, we will focus more on head and neck cancer issues which is a challenging horizon.

5. Source of Funding

None.

6. Conflict of Interest

None.

References

- Roesch KZ, Rocco J, Yom SS, Haddad R, Saba NF. Head and neck cancer. *Lancet*. 2022;398(10318):2289–98. doi:10.1016/S0140-6736(21)01550-6.
- Kohan EJ, Wirth WA. Anatomy of the neck. *Clin Plastic Surg*. 2014;41(1):1–6.
- Shadfar S, Perkins SW. Anatomy and physiology of the aging neck. *Facial Plast Surg Clin North Am*. 2014;22(2):161–70.
- Ihnatsenka B. Applied sonoanatomy of the posterior triangle of the neck. *Int J Shoulder Surg*. 2010;4(3):63–74.
- Guidera AK, Stringer MD. Head and neck fascia and compartments: no space for spaces. *Head Neck*. 2014;36(7):1058–68.
- Warshafsky D. Imaging anatomy of deep neck spaces. *Otolaryngol Clin North Am*. 2012;45(6):1203–21.
- Frisdal A, Trainor PA. Development and evolution of the pharyngeal apparatus. *Wiley Interdiscip Rev Dev Biol*. 2014;3(6):403–18. doi:10.1002/wdev.147.
- Garner DH. StatPearls Publishing; Treasure Island (FL): Anatomy, Head and Neck. *Carotid Sheath*. 2022;2(2):12–22.
- Lee JH. High-resolution Imaging of Neural Anatomy and Pathology of the Neck. *Korean J Radiol*. 2017;18(1):180–93.
- Ha EJ. Clinical significance of vagus nerve variation in radiofrequency ablation of thyroid nodules. *Eur Radiol*. 2011;21(10):2151–7.
- Masuoka H. Innervation of the cricothyroid muscle by the recurrent laryngeal nerve. *Head Neck*. 2016;38(1Suppl 1):441–5. doi:10.1002/hed.24015.
- Hwang K. Anatomy of the Platysma Muscle. *J Craniofac Surg*. 2017;28(2):539–42.
- Bordoni B, Varacallo M. Anatomy, Head and Neck, Sternocleidomastoid Muscle. vol. 1. StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022.
- Ourieff J, Scheckel B, Agarwal A. Anatomy, Back, Trapezius. vol. 3. StatPearls Publishing; Treasure Island (FL); 2022. p. 31–41.
- Tan BK, Wong CH, Chen HC. Anatomic variations in head and neck reconstruction. *Semin Plast Surg*. 2010;24(2):155–70. doi:10.1055/s-0030-1255333.

16. Falyar CR. Localization of the brachial plexus: Sonography versus anatomic landmarks. *J Clin Ultrasound*. 2016;44(7):411–5.
17. Cappiello J. The spinal accessory nerve in head and neck surgery. *Curr Opin Otolaryngol Head Neck Surg*. 2007;15(2):107–11.
18. Kim DH, Cho YJ, Tiel RL, Kline DG. Surgical outcomes of 111 spinal accessory nerve injuries. *Neurosurgery*. 2003;53(5):1102–12. doi:10.1227/01.neu.0000089058.82201.3d.
19. Allen E, Minutello K, Murcek BW. StatPearls Publishing; Treasure Island (FL): Anatomy, Head and Neck. *Larynx Recurrent Laryngeal Nerve*. 2022;2(2):25–35.
20. Chiba M. Clinical Features of Idiopathic Parotid Pain Triggered by the First Bite in Japanese Patients with Type 2 Diabetes: A Case Study of Nine Patients. *Pain Res Treat*. 2018;20(19):7861451. doi:10.1155/2018/7861451.
21. Cunha B, Tadi P, Bragg BN. Torticollis. vol. 12. StatPearls Publication, Treasure Island (FL):Torticollis; 2022. p. 129–39.
22. Richard M. Embryology, Anatomy, Histology, and Developmental Anomalies of the Liver. vol. 71. Sleisenger and Fordtran's Gastrointestinal and Liver Disease; 2022. p. 1129–34.

Author biography

V Rama Raju, Professor

Cite this article: Raju VR. Analysis of anatomy of head and neck: A study through empiricals part- II. *IP Indian J Anat Surg Head, Neck Brain* 2022;8(4):129-132.